

Stringent New Bounds on Supersymmetric Higgs Bosons from Existing Tevatron Data

Manuel Drees¹, Monoranjan Guchait^{2,3}, and Probir Roy³

¹*APCTP, 207-43 Cheongryangryi-dong, Tongdaemun-gu, Seoul 130-012, Korea*

²*Physics Department, Yonsei University, Seoul 120-749, Korea*

³*Tata Institute of Fundamental Research, Mumbai, India*

There is an error in our reported calculation [1] of the cross section for associated $b\bar{b}$ Higgs production in hadron colliders. After correction, the cross section at the Tevatron is reduced by a factor of about 10. (Our calculations have been performed by choosing the Higgs mass as the QCD scale and CTEQ3L structure functions; there could be an uncertainty of upto 40% from different choices). This weakens (increases) the upper bound on $\tan\beta$ that can be derived from the existing CDF data [2] by slightly more than a factor of 3. The excluded region, in the plane spanned by the MSSM parameters m_A and $\tan\beta$, is shown in Fig. 1, which corrects Fig. 2 of our original publication [1]. We are still able to extend the region that can be excluded from a consideration of $t \rightarrow H^+$ decays [3], though our upper bound on $\tan\beta$ is always above 80. However, a similar analysis, applied to the forthcoming data on $b\bar{b}\tau^+\tau^-$ final state events from RUN 2 at the Tevatron, would be able to lead to much stronger conclusions.

We thank Steve Mrenna for alerting us to the problem in our calculation, and Michael Spira for providing us with results of his independent calculation.

References

- [1] M. Drees, M. Guchait and P. Roy, hep-ph/9801229, Phys. Rev. Lett. **80**, 2047 (1998).
- [2] CDF collab., F. Abe et al., Phys. Rev. Lett. **78**, 2906 (1997).
- [3] M. Guchait and D.P. Roy, Phys. Rev. **D55**, 7263 (1997). See also CDF collab., F. Abe et al., Phys. Rev. Lett. **79**, 357 (1997).

Figure Captions

Fig. 1: Constraints on the MSSM Higgs sector in the $(m_A, \tan\beta)$ plane. The region $m_A < 75$ GeV is excluded by Higgs searches at the LEP collider. The region above the dashed curve is excluded by an analysis [2] of top quark decays. The region above the solid line is excluded by our analysis using CDF limits on $b\bar{b}\tau^+\tau^-$ final states. For $\tan\beta \gg 1$, the CP-odd scalar A is nearly degenerate with either the light CP-even scalar h or the heavy CP-even state H ; the cross-over occurs between 100 and 125 GeV, depending on the details of the sparticle spectrum.

